



Cambridge IGCSE™

COMBINED SCIENCE

0653/42

Paper 4 Theory (Extended)

February/March 2022

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the February/March 2022 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **10** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1	Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
2	The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
3	Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
4	The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
5	<p><u>'List rule' guidance</u></p> <p>For questions that require <i>n</i> responses (e.g. State two reasons ...):</p> <ul style="list-style-type: none"> • The response should be read as continuous prose, even when numbered answer spaces are provided. • Any response marked <i>ignore</i> in the mark scheme should not count towards <i>n</i>. • Incorrect responses should not be awarded credit but will still count towards <i>n</i>. • Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should not be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response. • Non-contradictory responses after the first <i>n</i> responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks												
1(a)(i)	banana tree → macaw → python ; <i>one mark for correct order</i> <i>one mark for arrows in correct direction</i>	2												
1(a)(ii)	banana eaten by monkey and/or parrot ; monkey and/or parrot eaten by jaguar ;	2												
1(b)	<table border="1"> <thead> <tr> <th>part of leaf</th> <th>feature</th> <th>function</th> </tr> </thead> <tbody> <tr> <td>palisade mesophyll layer</td> <td>(many) chloroplasts</td> <td>(photosynthesis)</td> </tr> <tr> <td>spongy mesophyll layer</td> <td>(air spaces)</td> <td>gas exchange</td> </tr> <tr> <td>xylem tissue</td> <td>(hollow)</td> <td>transport of water / support</td> </tr> </tbody> </table> <div style="text-align: right;">⋮</div>	part of leaf	feature	function	palisade mesophyll layer	(many) chloroplasts	(photosynthesis)	spongy mesophyll layer	(air spaces)	gas exchange	xylem tissue	(hollow)	transport of water / support	3
part of leaf	feature	function												
palisade mesophyll layer	(many) chloroplasts	(photosynthesis)												
spongy mesophyll layer	(air spaces)	gas exchange												
xylem tissue	(hollow)	transport of water / support												
1(c)	as light intensity increases the rate of photosynthesis increases ; then levels off / owtte ;	2												

Question	Answer	Marks
2(a)	Z and has 7 electrons in its outer shell / 7 valence electrons ;	1

Question	Answer	Marks															
2(b)(i)	X in Group 2 (no mark) X atoms have two outer electrons; X is a metal ;	2															
2(b)(ii)	(X is metallic and) Y is non-metallic because it is a gas / exists as single atoms / X is a metal and Y is a noble gas / X atoms bond together and Y atoms not (normally) bonded / owtte ;	1															
2(b)(iii)	helium ;	1															
2(b)(iv)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Statement</th> <th>True</th> <th>False</th> </tr> </thead> <tbody> <tr> <td>The bond between two atoms in Z₂ is a double bond.</td> <td></td> <td>✓</td> </tr> <tr> <td>The bonding in Z₂ is covalent.</td> <td>✓</td> <td></td> </tr> <tr> <td>Z₂ molecules are diatomic.</td> <td>✓</td> <td></td> </tr> <tr> <td>Z₂ is non-metallic.</td> <td>✓</td> <td></td> </tr> </tbody> </table> <p>All correct (2) 2 / 3 correct (1)</p>	Statement	True	False	The bond between two atoms in Z ₂ is a double bond.		✓	The bonding in Z ₂ is covalent.	✓		Z ₂ molecules are diatomic.	✓		Z ₂ is non-metallic.	✓		2
Statement	True	False															
The bond between two atoms in Z ₂ is a double bond.		✓															
The bonding in Z ₂ is covalent.	✓																
Z ₂ molecules are diatomic.	✓																
Z ₂ is non-metallic.	✓																
2(b)(v)	X ion positive and Z ion negative ; X ion is +2 and Z ion is –1; (subsumes first point) X loses (2) electrons and Z gains (1) electron ;	3															

Question	Answer	Marks
3(a)(i)	friction ;	1
3(a)(ii)	25 (N) ;	1

Question	Answer	Marks
3(b)	KE = $\frac{1}{2}mv^2$ (<i>in any form</i>) / $\frac{1}{2} \times 20 \times 0.7 \times 0.7$; 4.9 (J) ;	2
3(c)(i)	use of area under curve ; calculation of one area e.g., distance = $(\frac{1}{2} \times 1 \times 0.7 = 0.35)$ / $(3 \times 0.7 = 2.1)$; (= 0.35 + 2.1 =) 2.45 (m) ;	3
3(c)(ii)	acceleration = change in speed \div time (<i>in any form</i>) / $0.7 \div 1$; 0.7 ; m / s ² ;	3

Question	Answer	Marks
4(a)(i)	X – oesophagus ; Y – pancreas ;	2
4(a)(ii)	produce / secrete, <u>amylase</u> ; digest <u>starch</u> ; to produce, simpler sugars / glucose ;	3
4(b)	prevents constipation ;	1
4(c)	water moves from higher to lower water potential ; through a partially permeable membrane ; (by the process of) osmosis ;	3
4(d)	thin <u>wall</u> / <u>wall</u> one cell thick ; allow (efficient) exchange of substances ;	2

Question	Answer	Marks
5(a)	hydrogen: lighted splint and pops / gives a squeaky pop ; oxygen: glowing splint and relights ;	2
5(b)(i)	$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$ $4\text{OH}^-(\text{aq}) \rightarrow \text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^-$ (g) shown for hydrogen and oxygen ; (aq) for H^+ and OH^- and (l) for water ;	2
5(b)(ii)	(at the negative electrode / cathode) hydrogen ions gain electrons (to form hydrogen atoms / hydrogen molecules) ; (at the positive electrode / anode) hydroxide ions lose electrons (to form water molecules and oxygen molecules) ;	2
5(c)	universal indicator / (blue) litmus AND turns red ;	1

Question	Answer	Marks
6(a)	<u>focal length</u> ;	1
6(b)(i)	X located between A and lens ;	1
6(b)(ii)	density = mass \div volume (<i>in any form</i>) / $m = V \times \rho = 50.0 \times 0.85$; 42.5 (g) ;	2
6(c)(i)	kinetic energy of molecules increases ; distance between molecules increases / molecules further apart ;	2
6(c)(ii)	(decreases – no mark) higher volume but same mass / the idea that $m \div V$ decreases ;	1

Question	Answer	Marks
7(a)(i)	300 – 280 ; 20 ;	2
7(a)(ii)	tar ; oxygen ; nicotine ;	3
7(b)	(damaged alveoli) less surface (area) ; so less diffusion / lower rate of diffusion (of gases) ;	2

Question	Answer	Marks																				
8(a)(i)	cracking ;	1																				
8(a)(ii)	high temperature ; catalyst / high pressure ;	2																				
8(b)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th>C₇H₁₆</th> <th>C₂H₄</th> <th>C₅H₁₂</th> </tr> </thead> <tbody> <tr> <td>It has molecules which contain 23 atoms.</td> <td style="text-align: center;">✓</td> <td></td> <td></td> </tr> <tr> <td>It is unsaturated.</td> <td></td> <td style="text-align: center;">✓</td> <td></td> </tr> <tr> <td>It is an alkane.</td> <td style="text-align: center;">✓</td> <td></td> <td style="text-align: center;">✓</td> </tr> <tr> <td>It produces carbon dioxide and water when it burns.</td> <td style="text-align: center;">✓</td> <td style="text-align: center;">✓</td> <td style="text-align: center;">✓</td> </tr> </tbody> </table> <p>All correct (3 marks) 3 rows correct (2 marks) 2 rows correct (1 mark)</p>		C ₇ H ₁₆	C ₂ H ₄	C ₅ H ₁₂	It has molecules which contain 23 atoms.	✓			It is unsaturated.		✓		It is an alkane.	✓		✓	It produces carbon dioxide and water when it burns.	✓	✓	✓	3
	C ₇ H ₁₆	C ₂ H ₄	C ₅ H ₁₂																			
It has molecules which contain 23 atoms.	✓																					
It is unsaturated.		✓																				
It is an alkane.	✓		✓																			
It produces carbon dioxide and water when it burns.	✓	✓	✓																			
8(c)	<i>from orange to colourless ;</i>	1																				

Question	Answer	Marks
8(d)	(same) general formula ; similar chemical properties ;	2

Question	Answer	Marks														
9(a)	variable resistor ; controls current / the idea that it controls the brightness of the lamps ;	2														
9(b)	resistance of each lamp = $V \div I / 220 \div 5 / 44 (\Omega)$; combined resistance = $(R_1 \times R_2) \div (R_1 + R_2)$ (<i>in any form</i>) / $44 \times 44 \div 88$; $22 (\Omega)$;	3														
9(c)(i)	the current (through lamps) exceeded 10 A; because the current fluctuates / fuse needs to allow for fluctuations / fuse rating needs to be a little higher than expected current / owtte ;	2														
9(c)(ii)	the idea that a 13 A fuse allows normal operation / takes 10 A without blowing ; a rating greater than 13 A risks damaging the circuit / lamps / owtte ;	2														
9(d)	<table border="1" style="width: 100%; text-align: center;"> <tr> <td colspan="7">← increasing frequency</td> </tr> <tr> <td>(gamma radiation)</td> <td>(X-rays)</td> <td>ultraviolet (radiation)</td> <td>visible (light)</td> <td>infrared (radiation)</td> <td>(microwaves)</td> <td>(radio waves)</td> </tr> </table>	← increasing frequency							(gamma radiation)	(X-rays)	ultraviolet (radiation)	visible (light)	infrared (radiation)	(microwaves)	(radio waves)	1
← increasing frequency																
(gamma radiation)	(X-rays)	ultraviolet (radiation)	visible (light)	infrared (radiation)	(microwaves)	(radio waves)										